

US EPA ARCHIVE DOCUMENT

# TRACING NITROGEN MOVEMENT IN FORESTED WATERSHEDS: PRELIMINARY RESULTS FROM THE SLEEPERS RIVER RESEARCH WATERSHED, VT

Stephen D. Sebestyen<sup>1</sup>, Elizabeth W. Boyer<sup>2</sup>, James B. Shanley<sup>3</sup>, Nobuhito Ohte<sup>4</sup>, Daniel H. Doctor<sup>5</sup>, & Carol Kendall<sup>5</sup>


<sup>1</sup> State University of NY College of Environmental Science & Forestry, Syracuse, NY; <sup>2</sup> UC Berkeley, Berkeley, CA; <sup>3</sup> USGS, Montpelier, VT; <sup>4</sup> Kyoto University, Kyoto, Japan; <sup>5</sup> USGS, Menlo Park, CA



## INTRODUCTION


Human activities have increased (nitrogen) availability in the environment & stream export of dissolved N has increased in many regions

Our work assesses how combined hydrological & biogeochemical processes control variation of stream  $\text{NO}_3^-$  (nitrate) concentrations over multiple time scales (event, seasonal, & annual). We used high-frequency measurements (hydrological, chemical, & isotopic tracers) to fingerprint both water flow paths and N sources to the stream from the landscape.




## NITRATE ISOTOPES & SNOWMELT

- Stream  $\text{NO}_3^-$  concentrations were strongly influenced by atmospheric  $\text{NO}_3^-$  during early snowmelt
- During peak snowmelt,  $\text{NO}_3^-$  was from atmospheric & soil (microbially nitrified) sources
- In late snowmelt, stream  $\text{NO}_3^-$  was from soil & groundwater  $\text{NO}_3^-$  sources



## NITRATE & ANTECEDENT DISCHARGE

- $\text{NO}_3^-$  concentrations in the stream are highest when soil N concentrations are highest, highlighting the importance of terrestrial-aquatic linkages
- Soil  $\text{NO}_3^-$  is highest during the dormant season when biotic uptake is minimal & creates a pool of N that is available to flush to streams with snowmelt
- $\text{NO}_3^-$  is higher in soils & streamflow after dry antecedent moisture conditions than after wet



Cumulative flow of water at the W-9 weir was calculated for the 7 day interval prior to water sampling. Low cumulative flow indicates low water fluxes and dryer conditions prior to sample collection

## IMPLICATIONS

- Stream  $\text{NO}_3^-$  variation is influenced by hydrological flushing of source areas that vary with wetness
- Highest  $\text{NO}_3^-$  concentrations occur with the first pulses of high flow after dry periods, highlighting moisture controls
- During high flow, water moved rapidly to streams via preferential flowpaths (overland & shallow subsurface) and thereby shortcuts retention in the landscape
- Atmospheric deposition is an important N source to the landscape as reflected in stream chemistry and especially during early snowmelt when infiltrating water has a short residence time in the landscape

